

Seasonal Aerosol Classification Over the South Asia By Using Satellite-Based Atmospheric Optical Data

Anum Liaqut^{1*}, Isma Younes¹, Salman Tariq^{2,3}

¹Department of Geography, University of the Punjab, Lahore, Pakistan

²Remote sensing, GIS and Climatic Research Lab (National Center of GIS and Space Application, University of the Punjab, Lahore, Pakistan

³Department of Space Science, University of the Punjab, Lahore, Pakistan

* Email: arjandas99@gmail.com

Abstract: Aerosol optical characteristics have been investigated to explore regional and seasonal inconsistencies of aerosols and to define the dominant type throughout South Asia from 2001 to 2021. MODIS aerosol products from collection 6.1 have been used in present study, that comprise daily values of Angstrom exponent (AE) and aerosol optical depth (AOD) data. MODIS-derived AODs are validated by using nine ground-based AERONET station data. Overall, an adequate correlation is found among the two datasets. However, an overestimation of the MODIS retrievals is found in one site named Jaipur and underestimations are found in two sites named as Gandhi-college and Karachi. The seasonal evaluation shows that aerosol distribution found between 0 and 1.05, depending on the change in geographical location. The highest AOD value is originate over the Indo-Gangetic plain (IGP), mostly throughout warm season. The second maximum AOD value covers a large area of South Asia during spring, summer and autumn. The lowest values of AOD are found in the winter season excluding the IGP. A region with high aerosol optical depth (AOD) values are support a low value of angstrom exponent (AE) indicating the coarse aerosol during warm seasons (spring and summer) over IGP. The region with high AOD and high AE values is showing fine aerosol during the mild to cold seasons (autumn and winter). The threshold values for AOD and AE have been used to classify aerosols. The results demonstrate that urban/industrial aerosols prominent in every season across the region but dominate in spring and summer due to the frequent occurrence of dust events. The mixed type aerosol is the second largest contributor in aerosol formation in all seasons. The Biomass burning/smoke aerosol is dominant over IGP due to open forest and crop burning in autumn. Clean and maritime aerosol has small unnoticeable involvement in the studied region.

Keywords: Aerosol, classification, IGP, south asia.

Introduction

Aerosols are the most changeable element of the Earth's atmosphere that have been identified as a key contributor to the climate and radiative balance (Zhu, 2019). Aerosols can absorb or deflect land and solar radiation and producing the heating and cooling effect for earth system that base on their optical and physical characteristics (Giles, 2012). Hygroscopic aerosols can also contribute as condensation nuclei and affect cloud reflectance and rainfall formation in an indirect manner (Alizadeh, 2018). Different optical and microphysical properties of aerosol have been used to classify of aerosol around the world (Valentini, 2020). The AOD and AE are the two major columnar aerosol characteristics used to categorize aerosols. The first is used to calculate the amount of aerosol, whereas the second is an estimate of aerosol size. Mhawish et al. (2017) analyzed AOD retrieval algorithms for classification of aerosol, such as fine, mixed and coarse aerosols. Dust aerosols have a low AE and high AOD value, whereas maritime aerosols have a high AE and low AOD value. In contrast, Sreekanth (2014) identified the anthropogenic aerosols by using high values of Aerosol optical depth and Angstrom Exponent. The following is the most prevalent aerosol categorization in the literature: (a) desert dust; (b) industrial/urban aerosols released by burning of fossil

fuel; (c) maritime aerosols; (d) smoke/BB from forest fires and open crop burning and (e) mixed type of aerosols (Hamill, 2016). Due to the lack of a ground-based network, satellite network is the primary method for determining spatial and temporal variability of aerosol at local to global level (Sabetghadam, 2018). Due to significant uncertainties in satellite-derived aerosol products, ground-based aerosol data have been used to validate the satellite-based aerosol data (Benkhalifa, 2017). MODIS provides significant and good worldwide coverage, as well as nearly daily AOD data, which indicates aerosol load in the atmosphere that influence the radiation budget of earth (Luo, 2014). Numerous studies have conducted to compare and evaluate satellite-derived AOD with ground-based aerosol data (Rupakheti, 2018) across the world (Boiyo, 2017; Bilal, 2018a; 2018b; 2019; Liu, 2018). Nevertheless, the satellite-derived AOD is always with some anomalies because of the diverse landform, and cloud contamination (Kang, 2016).

In the present study, aerosol parameters AE and AOD are studied over South Asian region for the past two decades; from 2001 to 2021 to examine regional and seasonal variations of aerosols. Ground-based AERONET aerosol data has been used to validate the satellite-based aerosol data to verify the accuracy of satellite-based products. The satellite-based

Copyright © SEGMITE

classification technique, using AE and AOD is applied to examine the contribution of the leading type of aerosol over South Asia. The results are evaluated to provide an understanding of the spatiotemporal distribution and seasonal pattern of the region's aerosol types.

Material and Methods

Study area

South Asia contains the eight countries according to SAARC, named Pakistan, India, Bhutan, Nepal, Afghanistan, Bangladesh, Maldives and Sri Lanka. South Asia generally includes the land around the Arabian Sea (Fig 1). The region has three different climate zones. The climate condition of South Asia is highly affected by the pattern of the monsoon that brings the irregular phases of dry and wet weather. Summer (JJA) in South Asia experiences the wet monsoon due to the moist winds that blow from the sea, whereas due to the dry winds that come from land, winter (DJF) experiences the dry monsoon in the study region (Ramachandran et al., 2015). For the detailed study, highly polluted cities that are influenced by different anthropogenic emissions, along with natural

Different urban region in South Asia have different natural and man-made activities that produced different aerosol sources. Thus, aerosol properties in South Asia have been highly fluctuating on temporal and spatial scales (Lawrence and Lelieveld, 2010).




Fig. 1 Google earth-based Locations of selected AERONET sites.

Table 1 Details of AERONET sites.

Aeronet sites	Lat/Long	Elevation (m) AMSL	Time-period	Prominent Source of Aerosols
Bhola	22.2°N/90.7°E	7	2014-2018	Urban industries, Crop-residue burning, brick kilns, Mining activities, power plants.
Dhaka	23.7°N/90.3°E	34	2014-2018	Urban industries, Crop burning, power plants megacity.
Gandhi College	25.8°N/84.1°E	22	2006-2019	Crop burning in pre-monsoon and post-monsoon, Small and large industries, brick kilns industries, power plants, home heating, trans-boundary.
Jaipur	26.9°N/75.8°E	450	2011-2017	
Kanpur	26.5°N/80.2°E	123	2011-2018	
Karachi	24.9°N/67.1°E	49	2010-2016	fossil fuel/ biomass burning, industrial hub, Windblown of desert dust, sea salt.
Lahore	31.4°N/74.2°E	209	2010-2016	Crop burning in pre-monsoon and post-monsoon, industrial hub, home heating, brick making industries, dust storms, power plants.
Lumbini	27.4°N/83.2°E	110	2013-2018	Urban industries, Power plants, home heating.
Pokhara	28.1°N/83.9°E	800	2010-2018	Urban Industries, power plants, home heating, brick making industries, trans-boundary.

aerosols, have been selected (World Air Quality Report, 2018). According to Intergovernmental Panel on Climate Change (2014) South Asia is likely to be affected by global warming that increase the dry and hot days in this zone. South Asia has experienced the maximum increase in AOD since 2010 (Pozzer, 2015). With the growing urbanization, intense industrialization and high population, the aerosol load is continuously rising in the region, especially in IGP (Wang, 2014). In contrast, the extreme aerosol loading over South Asia has the potential to cause a great number of early mortalities, a high risk of different and sever health issues such as cancer (Behrooz, 2020).

Satellite dataset

The MODIS terra and aqua satellites are optical sensor instruments that detect the related parameters of aerosol globally. In the present study, MODIS aerosol products in Level 3, collection 6.1 with $1^\circ \times 1^\circ$ spatial resolution are used over a time span from January 2001 to December 2021, which includes the AE and AOD at 550 nm for South Asian region. The deep blue algorithm is utilized for the obtaining of aerosol data over the study area. The MODIS AOD was retrieved at a 550 nm of wavelength for the study region (Remer, 2008). Conversely, the Angstrom Exponen shows the

size distribution of aerosols in the atmosphere. Angstrom exponent values greater than 1 are associated with the surplus of fine particles, while angstrom exponent values less than 1 demonstrate the dominance of coarse aerosol in the atmosphere (Remer, 2008). To identify the leading aerosol type, AE and AOD combinations are commonly used (Rupakheti, 2019; Chen, 2020). Daily average values of AE (Angstrom Exponent) and AOD (Aerosol Optical Depth) over South Asia from the MODIS aqua satellite are utilized in this study. Data is obtained from the website: <http://giovanni.gsfc.nasa.gov/giovanni>. These data sets are utilized to examine the temporal and spatial aerosol parameters and to identify the dominant type of aerosol over South Asia. Averaged data from summer (JJA; June, July, and August), spring (MAM; March, April, and May), winter (DJF; December, January, and February), and autumn (SON; September, October, and November) were examined. Furthermore, the grouping of AE and AOD data was used to identify the prominent season-based aerosol types.

AERONET dataset

AERONET is a ground-based sky-radiometers and sun-photometers station that provides real-time aerosol and related parameter data across the world (Holben, 2001). Data is obtained from <http://aeronet.gsfc.nasa.gov/>. The AERONET provided data in three levels: the first is level 1.0 data that contains pre-screened data. The second type of data is level 1.5 that includes cloud-screened data (Dubovik, 2000). 3rd is level 2.0 data that covers quality assured and cloud screened data (Giles, 2019) and this data can be utilized for a precise measurement of aerosol parameters (Dubovik, 2002). AERONET provided level 2.0 AOD data are significantly accurate with 0.01% uncertainties (Sayer, 2014) and that is why this data is generally utilized for verify of satellite observations. In the present study, AERONET level 2 version 3 daily based data is used for validation of satellite data from nine stations across Southeast Asia. The location of the selected sites is shown in table 1 and figure 1. These sites have been selected because of their long-term data series in South Asia.

Verification of satellite AOD with AERONET AOD

The long-term seasonal variation of AE and AOD is investigated over the study region. The uncertainty in

satellite measurements can be found by comparing ground-based and satellite-derived aerosol data. In the present study, Nine AERONET sites named in Table 1, are utilized for the assessment of the satellite-based (MODIS) data over South Asia. It is important to change the MODIS AOD and AERONET values on the same wavelength to achieve the purpose. AERONET AE (440–870 nm) and AOD 500 nm are used to convert the AERONET AOD 500 nm at the wavelength of 550 nm (Bibi, 2015):

$$AOD_{550} = AOD_{500} (550/500)^{-AE}$$

Algorithm for Aerosol classification

Assessment of various aerosol types is a significant problem for the climatic researcher that may give the detailed evaluation of the impact of radiation on aerosol (Satheesh, 2005). Gobbi and Barnaba (2004) have introduced a very appropriate algorithm to determine the different classes of aerosol based on AOD and AE. MODIS satellite retrievals are utilized for the aerosol classification over South Asia, where air masses and atmospheric circulation also affect aerosol types (Valentini, 2020). In this study, aerosols have been categorized into six classes, such as urban, smoke/biomass burning, clean, desert dust, maritime and mixed type aerosols. AE and AOD threshold values have been used for aerosol classification that mentioned in Table 2. This classification based on a literature review (Higurashi, 2002; Barnaba, 2004; Pace, 2006; Kaskaoustis, 2007; Mulcahy, 2009; Pathak, 2012; Pawar, 2015; Penning, 2015; Alam, 2016; Sabetghadam, 2020). It is important to mention that different threshold values may change the occurrence rate of each type. Thus, there are no set standards for these critical thresholds. These are generally site-specific (Patel, 2017).

Table 2. Threshold for aerosol classification.

Aerosol type	AE	AOD
Urban	>0.8	>0.25
Biomass burning/smoke	>1.3	>0.4
Maritime	<0.9	<0.15
Desert dust	<0.6	>0.4
Clean	<1	<0.2
Mixed type	Remaining	Remaining

Results and Discussion

Validation of AODs

Table 3 Summary of statistical metrics between the AERONET and MODIS AOD at each site.

Sr.	Sites	Days	r	RMB%	RMSE	EE %within (EE)	EE %above (+EE)	EE %below (-EE)
1	Bhola	193	0.80	17.4	0.26	62.7	3.6	33.7
2	Dhaka	349	0.77	17.2	0.38	64.7	2.6	32.6
3	Gandhi-College	1162	0.32	-35.1	0.41	35.8	8.4	56.7
4	Jaipur	520	-0.04	25.6	0.44	35.6	43.0	21.3
5	Kanpur	2603	0.80	11.0	0.15	69.7	7.4	22.9
6	Karachi	1673	0.32	-31.9	0.23	31.7	7.5	60.7
7	Lahore	1583	0.78	4.0	0.18	68.1	20.8	11.1
8	Lumbini	111	0.68	17.9	0.39	43.2	12.6	44.1
9	Pokhara	912	0.54	-67.8	0.53	18.1	1.8	80.1




Fig 2: Scatter plots between AERONET and MODIS AOD at 550 nm (λ) at the selected sites over South Asia including (a) Bhola, (b) Dhaka, (c) Gandhi-College, (d) Jaipur, (e) Kanpur, (f) Karachi, (g) Lahore, (h) Lumbini and (i) Pokhara. The r value is significant (p value <0.01) for all sites. The dashed blue lines illustrate the identity line, the solid blue line depicts the lower and upper Estimated Error area and the red line indicate the regression line.


Before the validation of MODIS satellite AOD retrieval over South Asia, it is important to note that the sampling techniques of satellite data are relatively different from AERONET (Sayer, 2014). Nevertheless, it contributes to a rough evaluation of the satellite demonstration in the aerosol optical depth (AOD) retrieval algorithm. There are limited AERONET sites and only a few give long-term data series for study, which makes it challenging to properly assess the errors in satellite retrievals. Fig.2a to Fig.2i show the validation of the daily average value of AERONET AOD and MODIS AOD with the regression equation at nine selected sites. Table 3 shows the related statistical metrics at the respective sites. The results show the justified correlation between the AERONET

and MODIS AOD data almost all over the selected sites in South Asia. The correlation coefficient value is >0.6 at five sites named Bhola, Dhaka, Kanpur, Lahore and Lumbini, representing the good correlation between MODIS AOD and AERONET data and has a high % of retrievals within the EE 62.7%, 64.7%, 69.7%, 68.1% and 43.2% respectively. Variances in correlation coefficient may be caused by the various aerosol types that cover the study area. Furthermore, it seems that the satellite-based AOD is influenced by the land cover type (Holben, 2001). Conversely, the r-value is found to be <0.6 at Gandhi-College, Karachi and Pokhara due to the predominance of coarse aerosol because the adjacent area is covered with desert (Zia et al., 2017; Alam et al 2011a). The MODIS-based AOD

is significantly overestimated (RMB = 25.6%), with 43% of the data being above the EE (+EE) at Jaipur and underestimates (RMB = -35.1%, -31.9%) with 56.7%, 60.7% of the data below the EE (- EE) at Gandhi-College and Karachi respectively. Rupakheti, 2019 and Bilal, 2016 evidently mentioned in their studies, that high underestimated AOD values are generally produced as a result of errors in the aerosol technique utilized for aerosol inversion along with the effect of the planetary boundary layer on MODIS-

January 2001 to December 2021. AOD values ranging from 0 to 1.05 show the inconsistency of aerosol loadings over the study area as a result of diverse topography (Fig. 3). Seasonal evaluation (Fig. 3a to d) indicates the maximum values of aerosol loads over IGP generally found in summer, spring and autumn, whereas the winter season also shows the high value of AOD over the eastern side of IGP in South Asia. Afghanistan has the lowest value of aerosols in all seasons. This is constant with earlier researches that found high value of AODs over South Asia as a result

Seasonal patterns of AOD and AE over study area



based AOD.

Fig. 3 represents the seasonal pattern of AOD as to describe the amount of aerosol over South Asia from

of forest and crop-residue burning events during autumn and dust storms during the warm (summer and spring) season (Zia, 2017). Seasonal AOD may be inconsistent due to various seasonal phenomena for the

development and elimination of aerosols in the atmosphere, accompanied by air masses that transference aerosols from distant sources in individual seasons (Kaskaoutis, 2012). Dust events eventually increases over a major part of South Asia in spring and summer, as a result an increase in AOD in warm season (Alam et al., 2011c). Alam (2014) presented in their study that dust activities occurring due to the Thar and Thal Deserts in northeastern and southeastern Pakistan and India are more common in the warm season than in cold season. Moreover, aerosols are transferred or removed by rainfall and wind, which primarily reduce the values of AOD because of the clear sky in winter. The spatial distribution of AOD values indicates the highest and second highest values of AOD range between 0.65 to 1.05 primarily originate over the IGP in South Asia, and the third-highest AOD value lies between 0.44 to 0.64, almost covering the entire subcontinent during the spring, summer, and

autumn seasons (Fig. 3b, c, and d). This is similar to previous studies that IGP is recognized as a hot-spot region in South Asia with the maximum rate of dust events in the warm season and open crop burning in autumn (Zia, 2017; Alam, 2011c). The third maximum AOD value is visible throughout India and Pakistan, and it is especially noticeable in the spring, summer, and autumn.

The seasonal distribution of the averaged Angstrom Exponent values is shown in Fig 4 for the period of 2001-2021. The AE values lie between 0 to 1.76, indicating the extensive range of aerosol sizes over the South Asia. Seasonal AE variation indicates the areas with the maximum values of AOD in Fig. 3 have significantly lower AE values (i.e., coarse aerosols) over some parts of South Asia that are more noticeable in warm season (summer and spring). The loading of coarse aerosols is possibly released from the deserts in




Fig. 4 Seasonal pattern of Angstrom exponent (AE, 470–660 nm), Like Fig. 3

the region. Moreover, areas with higher values of AOD and with higher values of AE (Fig 3 and 4) covers a major part of South Asia and shows the predominance of fine aerosols as a result of open crop and forest burning in autumn and winter. The size parameter AE is usually found to be larger during open burning activities, indicating the high proportion of finer aerosols (Kumar, 2015; Kumar, 2018a). The IGP is a significant aerosol hotspot region with annual average value of AOD > 0.5 throughout the year (Kumar, 2018a). The aerosol load increases even more through open burning months, with AE > 1.2 and AOD values > 1.0 (Kaskaoutis et al., 2014). The concentration of finer aerosol increased over IGP toward the open-burning downwind areas, assisted by north-westerly airflow (Kumar et al., 2018a).

Classification of Aerosol over study region

Classification of aerosol over South Asia in four seasons based on the threshold values that are represented in Table 2 is shown in Figure 5. Aerosol types have the same spatial resolution that is $1^\circ \times 1^\circ$ as the actual satellite data, utilized for the analysis. Seasonal aerosol types reveal that mixed aerosol type is dominate over Afghanistan in every season. Industrial/Urban aerosol types cover a major portion of South Asia in all seasons (Fig. 5a and d). This result is similar to the aerosol classification where BB/smoke and urban aerosol types are prominent aerosol types in South Asia (Ramachandran, 2020).

In South Asia during winter (Fig. 5a) and spring (Fig. 5b), mixed type aerosol is the second-highest contributor in aerosol formation after urban aerosols. IGP is known as an industrial hub in South Asia that may cause urban industrial and BB/smoke aerosol over the IGP zone. Different anthropogenic activities may




Fig. 5 Seasonal aerosol classification over the South Asia averaged during 2001–2021.

cause urban/industrial aerosol over the entire study area, and crop-residue burning, particularly in India and Pakistan, is the main source of BB/smoke aerosol throughout the autumn and winter (Kumar et al., 2018). Aforementioned studies have also showed the excessive contribution of urban and BB/smoke aerosols over the study area (Azhar, 2019) that is also good demonstrated in the spatial distribution of AOD in Fig. 3. In their study, Alizadeh-Choo bari et al. (2016) showed the westward transport of dust from the desert land of India to eastern Pakistan. The occurrence of frequent dust storms during spring and summer, a small amount of desert dust aerosols is also dominated in some parts of India and some parts of southern Pakistan. The long-range transport of different fine aerosols such as local pollution and regional sea/maritime aerosol from the Arabian sea may lead to the mixed type of aerosols into the local region (Eck, 2008; Basart, 2009). During winter and spring (Fig. 5a and b), the western parts of Pakistan and the middle parts of India are dominated by mixed-type aerosol in all seasons but experience more load of mixed type of aerosol during winter and spring. During summer and autumn, the mixed type of aerosol is leading over some parts of the area, although the distribution of remaining types is comparable to detected in the other three seasons, although with a decreased spatial variation. However, clean and maritime aerosols have a small measurable influence on aerosol spreading over the whole region.

Conclusion

Aerosol parameters such as AE and AOD have been studied over South Asia to observe the regional and seasonal variation of aerosols and to classify the aerosol from 2001 to 2021. There is a rational correlation between the satellite-derived AODs and the AERONET-derived AODs. Nevertheless, underestimation of MODIS AOD retrievals is found on two sites, Gandhi College and Karachi, where high AERONET AOD values are identified and overestimated on one site, Jaipur. Seasonal AOD patterns have been studied over South Asia. The highest values of AOD are found over IGP in South Asia during spring, summer and autumn seasons, while the eastern side of IGP in the winter season has the highest value of AOD. Second and third maximum AOD values are found over the rest of Pakistan and India, while Afghanistan, Bangladesh, Nepal and Bhutan experience the lowest values of AOD during all seasons. This seasonal variability is because of different seasonal phenomena that occur in the region, such as dust storms in spring and summer and crop and forest burning in autumn. Seasonal averaged value of AE lies between 0 and 1.76 that indicate the variation in aerosol size over the region. High values of Angstrom Exponent are found in autumn and winter over the eastern side of the region depict the dominance of fine aerosol because of crop burning and biomass burning, while low values of AE in summer indicate the high amount of coarse aerosol due to

desert dust. The seasonal aerosol classification demonstrates the high proportion of urban/industrial aerosol type during each season that almost cover the entire region. The mixed aerosol is the second-largest contributor in aerosol composition in each season, particularly in spring and winter. IGP experienced the biomass burning/smoke aerosol during autumn and winters. Afghanistan regularly experiences mixed aerosol types during the whole year. Lastly, clean and maritime aerosols have less noticeable involvement in the aerosol loading of the entire region.

References

- Azhar, R., Zeeshan, M., Fatima, K. (2019). Crop residue open field burning in Pakistan; multi-year high spatial resolution emission inventory for 2000–2014. *Atmos. Environ.*, **208**, 20-33.
- Alam, K., Shaheen, K., Blaschke, T., Chishtie, F., Khan, H.U., Haq, B.S. (2016). Classification of aerosols in an urban environment on the basis of optical measurements. *Aerosol Air Qual. Res.*, **16**(10), 2535-2549.
- Alam, K., Trautmann, T. and Blaschke, T. (2011a). Aerosol Optical Properties and Radiative Forcing over Mega-City Karachi. *Atmos. Res.*, **101**, 773-782.
- Alam, K., Qureshi, S. and Blaschke, T. (2011b). Monitoring Spatio-Temporal Aerosol Patterns over Pakistan Based on MODIS, TOMS and MISR Satellite Data and a HYSPLIT Model. *Atmos. Environ.*, **45**, 4641-4651.
- Alizadeh-Chooabari, O., Bidokhti, A.A., Ghafarian, P., Najafi, M.S. (2016). Temporal and spatial variations of particulate matter and gaseous pollutants in the urban area of Tehran. *Atmos. Environ.*, **141**, 443-453.
- Bilal M et al. (2021). Air pollution scenario over Pakistan: Characterization and ranking of extremely polluted cities using long-term concentrations of aerosols and trace gases. *Remo. Sens. Environ.*, **264**, 112-617.
- Behrooz, R.D., Kaskaoutis, D.G., Grivas, G., Mihalopoulos, N. (2020). Human health risk assessment for toxic elements in the extreme ambient dust conditions observed in Sistan, Iran. *Chemos.*, **262**, 127-835.
- Bibi, H., Alam, K., Chishtie, F., Bibi, S., Shahid, I., Blaschke, T. (2015). Intercomparison of MODIS, MISR, OMI, and CALIPSO aerosol optical depth retrievals for four locations on the Indo-Gangetic plains and validation against AERONET data. *Atmos. Environ.*, **111**, 113-126.
- Benkhalifa, J., L'eon, J.F., Chaabane, M. (2017). Aerosol optical properties of Western

- Mediterranean basin from multi-year AERONET data. *J. Atmos. Sol. Terr. Phys.*, **164**, 222-228.
- Boiyo, R., Kumar, K. R., & Zhao, T. (2017). Statistical intercomparison and validation of multisensory aerosol optical depth retrievals over three AERONET sites in Kenya, East Africa. *Atmos. Res.*, **197**, 277-288.
- Bilal, M., Nazeer, M., Qiu, Z., Ding, X., Wei, J. (2018a). Global Validation of MODIS C6 and C6.1 Merged Aerosol Products over Diverse Vegetated Surfaces. *Remote Sens.*, **10**(475), 1-17.
- Bilal, M., Qiu, Z., Campbell, J.R., Spak, S.N., Shen, X., Nazeer, M. (2018b). A New MODIS C6 Dark Target and Deep Blue Merged Aerosol Product on a 3 km Spatial Grid. *Remote Sens.*, **10**(463).
- Chen, Q.X., Huang, C.L., Yuan, Y., Mao, Q.J., Tan, H.P. (2020). Spatiotemporal distribution of major aerosol types over China based on MODIS products between 2008 and 2017. *Atmos.*, **11**(7), 703.
- Eck, T.F., Holben, B.N., Reid, J.S., Sinyuk, A., Dubovik, O., Smirnov, A., Giles, D., O'Neill, N.T., Tsay, S.C., Ji, Q., Al Mandoos, A. (2008). Spatial and temporal variability of column-integrated aerosol optical properties in the southern Arabian Gulf and United Arab Emirates in summer. *J. Geophys. Res. Atmos.*, **113** (D1).
- Giles, D.M., Sinyuk, A., Sorokin, M.G., Schafer, J.S., Smirnov, A., Slutsker, I., Eck, T.F., Holben, B.N., Lewis, J.R., Campbell, J.R., Welton, E.J., Korkin, S.V., Lyapustin, A.I. (2019). Advancements in the Aerosol Robotic Network (AERONET) Version 3 database Automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements. *Atmos. Meas. Tech.*, **12**, 169-209.
- Hamill, P., Giordano, M., Ward, C., Giles, D., Holben, B. (2016). An AERONET-based aerosol classification using the Mahalanobis distance. *Atmos. Environ.*, **140**, 213-233.
- Holben, B.N., Tanre, D., Smirnov, A., Eck, T.F., Slutsker, I., Abuhassan, N., Newcomb, W. W., Schafer, J.S., Chatenet, B., Lavenue, F., Kaufman, Y.J. (2001). An emerging ground-based aerosol climatology: aerosol optical depth from AERONET. *J. Geophys. Res. Atmos.*, **106**(D11), 12067-12097.
- IPCC: Intergovernmental Panel on Climate Change (2014) Climate Change 2013 – the Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
- Joshi, R. M. (2015). Education in South Asia. In: International Encyclopedia of the Social and Behavioral Sciences, **2nd edn**, 194-197.
- Kang, N., Kumar, K. R., Hu, K., Yu, X., & Yin, Y. (2016). Long-term (2002–2014) evolution and trend in Collection 5.1 Level-2 aerosol products derived from the MODIS and MISR sensors over the Chinese Yangtze River Delta. *Atmos. Res.*, **181**, 29-43.
- Kumar, M., Parmar, K.S., Kumar, D.B., Mhawish, A., Broday, D.M., Mall, R.K., Banerjee, T. (2018). Long-term aerosol climatology over Indo-Gangetic plain: trend, prediction and potential source fields. *Atmos. Environ.*, **180**, 37-50.
- Kumar, M., Tiwari, S., Murari, V., Singh, A.K., Banerjee, T. (2015). Wintertime characteristics of aerosols at middle Indo-Gangetic plain: impacts of regional meteorology and long-range transport. *Atmos. Environ.*, **104**, 162-175.
- Kumar, M., Ojha N., Singh N. (2021). Atmospheric aerosols from open burning in South and Southeast Asia. *Asian Atmospheric Pollution. Atmos. Environ.*, **106**, 168-181.
- Kumar, M., Parmar, K.S., Kumar, D.B., Mhawish, A., Broday, D.M., Mall, R.K., Banerjee, T. (2018). Long-term aerosol climatology over Indo-Gangetic Plain: trend, prediction and potential source fields. *Atmos. Environ.*, **180**, 37-50.
- Khoshshima, M., Bidokhti, A.A., Ahmadi-Givi, F. (2014b). Variations of aerosol optical depth and Angstrom parameters at a suburban location in Iran during 2009–2010. *J. Earth Syst. Sci.*, **123**(1), 187-199.
- Lawrence, M. G., Lelieveld, J. (2010). Atmospheric pollutant outflow from southern Asia: a review. *Atmos. Chem. Phys.*, **10**, 11017-11096.
- Liu, Y., Franklin, M., Kahn, R., Koutrakis, P. (2017). Using aerosol optical thickness to predict ground-level PM_{2.5} concentrations in the St. Louis area: a comparison between MISR and MODIS. *Remote Sens. Environ.*, **107**(1–2), 33-44.
- Mhawish, A., Banerjee, T., Sorek-Hamer, M., Bilal, M., Lyapustin, A.I., Chatfield, R., Broday, D.M. (2020). Estimation of high-resolution PM_{2.5} over the Indo-Gangetic plain by fusion of satellite data, meteorology, and land use variables. *Environ. Sci. Technol.*, **54**, 7891-7900.
- Patel, P.N., Dumka, U.C., Kaskaoutis, D.G., Babu, K.N., Mathur, A.K. (2017). Optical and radiative properties of aerosols over Desalpar, a remote site in western India: source identification, modification processes and aerosol type discrimination. *Sci. Total Environ.*, **575**, 612-627.
- Pathak, B., Bhuyan, P.K., Gogoi, M., Bhuyan, K., 2012. Seasonal heterogeneity in aerosol types over Dibrugarh-North-Eastern India. *Atmos. Environ.*, **47**, 307–315.

- Penning de Vries, M.J.M., Beirle, S., Hormann, C., Kaiser, J.W., Stammes, P., Tilstra, L.G., Wagner, T. (2015). A global aerosol classification algorithm incorporating multiple satellite data sets of aerosol and trace gas abundances. *Atmos. Chem. Phys.*, **15**(18), 10597-10618.
- Pozzer, A., De Meij, A., Yoon, J., Tost, H., Georgoulias, A.K., Astitha, M. (2015). AOD trends during 2001–2010 from observations and model simulations. *Atmos. Chem. Phys.*, **15**(10), 5521-5535.
- Ramachandran S., Rupakheti M. (2020). Year-round aerosol characteristics and radiative effects in the South Asian pollution outflow over a background site in the Maldives. *Atmos. Environ.*, **240**, 117-813.
- Rupakheti, D., Kang, S., Bilal, M., Gong, J., Xia, X., Cong, Z. (2019). Aerosol optical depth climatology over Central Asian countries based on Aqua-MODIS Collection 6.1 data: aerosol variations and sources. *Atmos. Environ.*, **207**, 205-214.
- Ramachandran, S., Kedia, S. V. (2015). Spatiotemporal characteristics of aerosols in India: observations and model simulations. *Atmos. Environ.*, **116**, 225-244.
- Rupakheti, D., S. Kang, M. Rupakheti, Z. Cong, L. Tripathi, A. Panday, and B. Holben. (2018). Observation of Optical Properties and Sources of Aerosols at Buddha's Birthplace, Lumbini, Nepal. *Environ. Sci. Poll. Res.*, **25**(15), 14868-14881.
- Saikawa, E., Wu, Q., Zhong, M., Avramov, A., Ram, K., Stone, E.A., Stockwell, C., Jayarathne, T., Panday, A., Yokelson, R. (2020). Garbage burning in South Asia – How important is it to regional air quality? *Environ. Sci. Technol.*, 341-349.
- Sreekanth, V. (2014). On the classification and sub-classification of aerosol key types over south central peninsular India: MODIS–OMI algorithm. *Sci. Total. Environ.*, **468**, 1086-1092.
- Sayer, A.M., Munchak, L.A., Hsu, N.C., Levy, R.C., Bettenhausen, C., Jeong, M.J. (2014). MODIS Collection 6 aerosol products: comparison between Aqua's e-Deep Blue, Dark Target, and "merged" data sets, and usage recommendations. *J. Geophys. Res. Atmos.*, **119**(24), 13-965.
- Sinyuk, A., Holben, B.N., Smirnov, A., Eck, T.F., Slutsker, I., Schafer, J.S., Giles, D.M., Sorokin, M. (2012). Assessment of error in aerosol optical depth measured by AERONET due to aerosol forward scattering. *Geophys. Res. Lett.*, **39**, 2-6.
- Sabetghadam, S., Khoshshima, M., Pierleoni, A. (2020). Aerosol climatology and determination of different types over the semi-arid urban area of Tehran, Iran: application of multi-platform remote sensing satellite data. *Atmos. Pollut. Res.*, **11**(9), 1625-1636.
- Valentini, S., Barnaba, F., Bernardoni, V., Calzolari, G., Costabile, F., Di Liberto, L., Forello, A.C., Gobbi, G.P., Gualtieri, M., Lucarelli, F., Nava, S. (2020). Classifying aerosol particles through the combination of optical and physical-chemical properties: results from a wintertime campaign in Rome (Italy). *Atmos. Res.*, **235**, 104-799.
- Wang, S., Fang, L., Gu, X., Yu, T., Gao, J. (2011). Comparison of aerosol optical properties from Beijing and Kanpur. *Atmos. Environ.*, **45**, 7406-7414.
- Zia H., Salman T., Ali M. (2017). Spatiotemporal patterns of correlation between atmospheric nitrogen dioxide and aerosols over South Asia. *Meteorol. Atmos. Phys.*, **129**, 507-527.
- Zhu, J., Xia, X., Che, H., Wang, J., Cong, Z., Zhao, T., Kang, S., Zhang, X., Yu, X., Zhang, Y. (2019). Spatiotemporal variation of aerosol and potential long-range transport impact over the Tibetan Plateau, China. *Atmos. Chem. Phys.*, **19**(23), 14637-14656.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).